

Factsheet: Laws of indices

Tom Coleman

Summary

A list of laws of indices.

The associated study guide for this factsheet is [Guide: Laws of indices](#). If you would like to know more about these, please read the guide.

Throughout this factsheet, a, b are positive real numbers, r and s are real numbers, and n is a positive whole number.

Laws of indices

Law 1: $a^r \cdot a^s = a^{r+s}$.

Law 2: $\frac{a^r}{a^s} = a^{r-s}$.

Law 3: $(a^r)^s = a^{r \cdot s}$.

Law 4: If a is non-zero, then $a^0 = 1$.

Law 5: $a^{-r} = \frac{1}{a^r}$

Law 6: $a^{1/n} = \sqrt[n]{a}$.

Law 7: $a^r \cdot b^r = (ab)^r$.

Law 8: $\frac{a^r}{b^r} = \left(\frac{a}{b}\right)^r$.

Law 9: $a^{1/n} \cdot b = \sqrt[n]{a \cdot b^n}$.

Law 10: $\sqrt[n]{a \cdot b} = \sqrt[n]{a} \cdot \sqrt[n]{b}$.

For more on the subject, please go to [Guide: Laws of indices](#)

Version history and licensing

v1.0: created in 05/24 by tdhc.

This work is licensed under [CC BY-NC-SA 4.0](#).